



European Society of Clinical Microbiology and Infectious Diseases
... improving the management of infections in Europe and beyond

RESEARCH NOTE

Corresponding author and reprint requests: N. S. Al-Waili, Life Support Technologies, Inc.,

**The Mount Vernon Hospital, 7th Avenue
12 North, Mount Vernon, NY 10550, USA**

E-mail: noori786@yahoo.com

Clinical and mycological benefits of topical application of honey, olive oil and beeswax in diaper dermatitis N. S. Al-Waili

Dubai Specialized Medical Center, Islamic Establishment for Education, Dubai, United Arab Emirates

ABSTRACT

Twelve infants suffering from diaper dermatitis were treated four times daily for 7 days with a mixture containing honey, olive oil and beeswax. The severity of erythema was evaluated on a five point scale. Three infants had severe erythema and ulceration, four had moderate erythema, and five had moderate erythema with maceration. The initial mean lesion score of 2.91 ± 0.79 declined significantly ($p < 0.05$) to 2.0 ± 0.98 (day 3), 1.25 ± 0.96 (day 5) and 0.66 ± 0.98 (day 7). *Candida albicans* was isolated initially from four patients, but from only two patients after treatment. This topical treatment was safe and well-tolerated, and demonstrated clinical and mycological benefits in the treatment of diaper dermatitis.

Continued ...

Keywords Beeswax, *Candida albicans*, dermatitis, diapers, honey, olive oil

Original Submission: 19 January 2004; Revised Submission: 16 July 2004; Accepted: 17 August 2004

Clin Microbiol Infect 2005; 11: 160–163
10.1111/j.1469-0691.2004.01013.x

160 Clinical Microbiology and Infection, Volume 11 Number 2, February 2005

2005 Copyright by the European Society of Clinical Microbiology and Infectious Diseases, CMI, 11, 145–163



European Society of Clinical Microbiology and Infectious Diseases
... improving the management of infections in Europe and beyond

Diaper dermatitis is the most common dermatological problem of infancy [1], occurring in 25–65% of children [2], and is caused by the combined irritant effects of wearing a diaper, urine and faeces [3]. Colonisation by *Candida* spp. is significantly more frequent in children with diaper dermatitis than in those with healthy skin, whereas colonisation by *Staphylococcus aureus* does not differ between these two groups [4]. Corticosteroids, zinc paste and eosin are well-known topical agents for the treatment of diaper dermatitis. It has been suggested that topical antifungal agents are not indicated for diaper dermatitis; indeed, their safety and effectiveness have not been established in infants [5].

Olive oil, beeswax and honey are natural products, containing flavonoids, and antioxidant, antibacterial and antifungal compounds, that affect the production of cytokines by skin cells when applied topically [6–11]. Previous studies have demonstrated the efficacy of a mixture containing honey, olive oil and beeswax (in a ratio of 1:1:1 v/v) for the treatment of dermatitis, psoriasis and skin fungal infections [12,13]. In the present pilot study, the effect of using this honey mixture to treat infants with diaper dermatitis was investigated.

Eight boys and four girls, aged 3–18 months and suffering from diaper dermatitis, were selected randomly for a pilot study. The infants developed dermatitis despite immediate bathing with warm water and replacement of a diaper following urination or defaecation. Informed consent was obtained from parents. At baseline, a thorough medical history was obtained and a complete physical examination was conducted. A moistened swab was used to sample the rash, erosion or ulceration, for mycological culture on Sabouraud glucose agar and subsequent testing for *Candida albicans*. Sampling was performed immediately before and at the end of treatment. Rash severity was assessed on a five-point scale (none = 0; mild erythema = 1; moderate erythema = 2; moderate erythema plus maceration = 3; and severe erythema plus pustules or ulceration = 4) at baseline and at days 3, 5 and 7 during treatment. Any new sign or symptom that appeared during therapy was recorded as an adverse effect. A positive therapeutic effect was recorded when a severe or moderate rash became mild or disappeared.

The topical treatment was prepared by thoroughly mixing natural honey, olive oil and beeswax (1:1:1 v/v; equivalent to (w/v) honey 50%, olive oil 29%, beeswax 21%). The honey was dark yellow in colour, and of multifloral origin; it contained (/100 g); 38 g fructose, 28 g glucose, water 20% v/v, 2.3 g vitamin C, 0.098 mg copper, 0.6 mg zinc, < 0.5 g sucrose, and 0.51 mg glutathione reductase. Natural olive oil prepared with the cold press method was used. The mixture was stored in the dark at room temperature until use.

The parents were asked to apply the topical treatment four times daily with gentle rubbing for a maximum of 7 days, and to bathe the infant with warm water and change the diaper following urination or defaecation. All parents used the same brand of diapers before and during the study. Infants who did not respond within 7 days were given conventional therapy, and the failure of the topical treatment was recorded. No other topical product was used during the study. The study was open, and it was deemed unethical to use a placebo on infants. Furthermore, any comparison with the use of standard therapy was postponed until the results of this pilot study were evaluated. Lesion scores were expressed as means \pm standard deviations. ANOVA tests were used for the comparison of mean scores before, during and after treatment, with $p < 0.05$ deemed to be significant. Culture of specimens collected from the lesions showed that *C. albicans* was present in four patients at baseline. *C. albicans* was associated with increased severity of rash. Of three infants who had severe erythema and ulceration, two were positive for *C. albicans*. Nine infants had moderate erythema, including five with maceration. The mean total rash score was 2.91 ± 0.79 at baseline (Table 1), decreasing to 2.0 ± 0.96



European Society of Clinical Microbiology and Infectious Diseases
... improving the management of infections in Europe and beyond

Table 1. Clinical and mycological effects of the topical treatment in patients with diaper dermatitis

Patient	Before treatment		After commencement of treatment			
	Lesion score	Culture for <i>Candida albicans</i>	Lesion score			Culture for <i>Candida albicans</i>
			3 days	5 days	7 days	
1	3	-	2	1	1	-
2	2	-	1	1	0	-
3	2	-	1	0	1	-
4	2	-	1	1	0	-
5	3	-	2	1	0	-
6	2	-	1	0	0	-
7	4	+	3	2	2	+
8	4	+	3	3	3	+
9	4	+	3	2	1	-
10	3	+	2	1	0	-
11	3	-	2	1	1	-
12	3	-	2	1	0	-
Mean ± SD	2.91 ± 0.79		2 ± 0.98 ^a	1.25 ± 0.96 ^a	0.66 ± 0.98 ^a	

^ap < 0.05.

(p < 0.05) at day 3 and to 1.25 ± 0.96 (p < 0.05) at day 5. At the end of the study, the mean total score was 0.66 ± 0.98, and ten of the 12 patients had either mild or no diaper dermatitis. *C. albicans* was eradicated from two of the four positive patients by the end of therapy. The two patients who showed no response to the study treatment had moderate erythema and were positive for *C. albicans*; these infants were treated with conventional therapy. No adverse effects were recorded, and the parents reported that the treatment was easy to apply and well-tolerated. The study treatment was effective in reducing the symptoms of diaper dermatitis, and eradicated *C. albicans* from 50% of culture-positive patients during the 7-day trial. Longer courses of treatment or more frequent topical applications might give even better results. *C. albicans* may not be the main causative agent of diaper dermatitis, and it is possible that bacteria could be involved in the aggravation of symptoms. However, patients who were culture-positive for *C. albicans* seemed to have aggravated symptoms of diaper dermatitis. The symptomatic improvement could be associated with the possible anti-inflammatory effects of the mixture, based on the properties of the ingredients. The mechanism of the antimicrobial action of honey has not been fully-defined, although acidity, osmolality and hydrogen peroxide production have been proposed as important factors [14]. Honey has been shown to increase nitric oxide levels and decrease prostaglandin levels in biological fluids [15–17]. A diet rich in olive oil has been shown to increase nitric oxide and decrease

arachidonic acid production by resident rat macrophages [18]. Nitric oxide released from the skin through the sweat glands has antimicrobial activity [19,20]. Therefore, the effects of the topical treatment might be caused by a reduction in prostaglandin synthesis at the site of application, elevation of nitric oxide concentrations in the lesions, inhibition of fungal or bacterial growth, and antioxidant and anti-inflammatory activities [6,7,11]. Considering the potential hazards of topical corticosteroids, it seems that use of this topical treatment, alone or in combination with other agents, is a possible alternative therapy for diaper dermatitis. This could result in reduced use of prescription drugs and a concomitant reduction in any associated adverse effects.

REFERENCES

- Hurwitz S. Clinical pediatric dermatology, 1st edn. Philadelphia: Saunders, 1981; 27–30.
- Jodan W, Lawson K, Berg R, Franxman J, Marrer A. Diaper dermatitis: frequency and severity among a general infant population. *Pediatr Dermatol* 1986; 3: 198–207.
- Atherton J. The aetiology and management of irritant diaper dermatitis. *J Eur Acad Dermatol Venerol* 2001; 15: 1–4.
- Ferrazzini G, Kaiser R, Hirsig Chen K, Wehrli M. Microbiological aspects of diaper dermatitis. *Dermatology* 2003; 206: 136–141.
- Hoppe E. Treatment of oropharyngeal candidiasis and candidal diaper dermatitis in neonates and infants: review and reappraisal. *Pediatr Infect Dis J* 1997; 16: 885–894.
- Tuck K, Hayball P. Major phenolic compounds in olive oil. *J Nutr Biochem* 2002; 13: 636–644.
- Noa M, Mas R. Effect of D-002 on the pre-ulcerative phase of carrageenan-induced colonic ulceration in the guinea pig. *J Pharm Pharmacol* 1998; 50: 549–553.
- Al-Waili N, Saloom K. Effects of topical honey on postoperative wound infections due to gram positive and gram negative bacteria following caesarean sections and hysterectomies. *Eur J Med Res* 1999; 26: 126–130.
- Al-Waili N. Investigating the antimicrobial activity of natural honey and its effects on the pathogenic bacterial infections of surgical wounds and conjunctiva. *J Med Food* 2004; 7: 210–222.
- Al-Waili N. Therapeutic and prophylactic effects of crude honey on chronic seborrheic dermatitis and dandruff. *Eur J Med Res* 2001; 6: 306–308.



European Society of Clinical Microbiology and Infectious Diseases
... improving the management of infections in Europe and beyond

11. Tranter S, Tassou C, Nychas J. The effect of the olive phenolic compound, oleuropein, on growth and enterotoxin B production by *Staphylococcus aureus*. *J Appl Bacteriol* 1993; 74: 253–259.
12. Al-Waili N. Topical application of natural honey, beeswax and olive oil mixture to treat patients with atopic dermatitis or psoriasis: partially controlled study. *Complement Ther Med* 2003; 11: 226–222.
13. Al-Waili N. Safety and efficacy of topical application of honey, olive oil and beeswax mixture in patients with tinea versicolor, tinea cruris, tinea corporis and tinea faciei. Pilot study. *Complement Ther Med* 2004; 12: 45–47.
14. Molan P. The role of honey in the management of wounds. *J Wound Care* 1999; 8: 415–418.
15. Al-Waili N, Boni N. Effects of honey ingestion on nitric oxide in saliva. *FASEB J* 2003; 17: 235.
16. Al-Waili N. Identification of nitric oxide metabolites in various honeys and effects of honey on plasma and urinary nitrite/nitrate concentration. *J Med Food* 2003; 6: 350–357.
17. Al-Waili N. Honey lowers plasma prostaglandin concentrations in normal individuals. *J Med Food* 2003; 6: 129–133.
18. Visioli F, Bellosta S, Galli C. Oleuropein. The bitter principle of olives enhances nitric oxide production by mouse macrophages. *Life Sci* 1998; 62: 541–546.
19. Al-Waili N. Identification of nitric oxide in sweat and tear. *FASEB J* 2003; 17: 342.
20. Akh U, Zefirov L. The physiological role of nitric oxide. *Usp Fiziol Nauk* 1999; 30: 54–72

Clinical Microbiology and Infection, Volume 11 Number 2, February 2005

2005 Copyright by the European Society of Clinical Microbiology and Infectious Diseases, CMI, 11, 145–163

